INTERNATIONAL
OLIVE OIL
COUNCIL

COI/T.15/NC no. 2/Rev. 8 25 November 1998

ENGLISH

Original: FRENCH

TRADE STANDARD APPLYING TO OLIVE OIL AND OLIVE-POMACE OIL

OLIVE OIL COUNCIL

RESOLUTION No. RES-3/79-IV/98

TRADE STANDARD APPLYING TO OLIVE OIL AND OLIVE-POMACE OIL

THE INTERNATIONAL OLIVE OIL COUNCIL,

Having regard to Resolution no. RES-2/78-IV/98 of 20 November 1997 whereby the IOOC adopted the trade standard applying to olive oil and olive-pomace oil, COI/T.15/NC no. 2/Rev. 7, which amended the standard COI/T.15/NC no. 2/Rev. 6 of 5 June 1997 as regards the maximum brassicasterol limit for olive-pomace oils (raised to 0.2%) and the specification that the trans fatty acid limit (C18:1T and C18:2T + C18:3T) must be less than or equal to the limit fixed for edible virgin olive oils;

Whereas the IOOC decided at its 79th session to raise the limit for linolenic acid content from 0.9% to 1.0%; to delete the possibility provided for in section 10.1.6.2. of declaring on the label the appellation of origin of the extra virgin olive oil used in blends with refined olive oil; to revise section 11 of the standard in order to delete the compulsory alkaline neutralisation of lampante virgin olive oil and crude olive-pomace oil prior to the determinations of the purity criteria; to insert ISO method 5555 "Animal and vegetable fats and oils – Sampling", ISO method 661 "Animal and vegetable fats and oils – Preparation of the test portion", IUPAC method no. 2.401 "Determination of the unsaponifiable matter" (in place of method COI/T.20/Doc. no. 10-5.1), and COI/T.20/Doc. no. 20 "Determination of the difference between real and theoretical ECN 42 triglyceride content" (in place of IUPAC method no. 2.324 and method COI/T.20/Doc. no. 9);

DECIDES

The trade standard applying to olive oil and olive-pomace oil, COI/T.15/NC no. 2/Rev. 8 of 25 November 1998 shall replace and rescind the trade standard applying to olive oil and olive-pomace oil COI/T.15/NC no. 2/Rev. 7 of 20 November 1997.

The Members shall take whatever measures are appropriate, in the manner required by their legislation, to apply the standard adopted and shall notify the Executive Secretariat of any such measures as soon as they are taken.

The non-Member States involved in international trade in olive oils and olive pomace-oils shall be invited to take into consideration the standard adopted and to adapt their regulations to the provisions thereof.

Florence (Italy), 25 November 1998.

OLIVE OIL

COUNCIL

COI/T.15/NC no. 2/Rev. 8 25 November 1998

ENGLISH

Original: FRENCH

TRADE STANDARD APPLYING TO OLIVE OIL AND OLIVE-POMACE OIL

1. SCOPE

This standard applies to olive oil and olive-pomace oil that is the object of international trade or of concessional or food aid transactions.

2. <u>DESIGNATIONS AND DEFINITIONS</u>

- **2.1.** Olive oil is the oil obtained solely from the fruit of the olive tree (Olea europaea sativa Hoffm. & Link), to the exclusion of oils obtained using solvents or re-esterification processes and of any mixture with oils of other kinds. It is marketed in accordance with the following designations and definitions:
- **2.1.1.** <u>Virgin olive oil</u> is the oil obtained from the fruit of the olive tree solely by mechanical or other physical means under conditions, particularly thermal conditions, that do not lead to alterations in the oil, and which has not undergone any treatment other than washing, decantation, centrifugation and filtration.

2.1.1.1. Virgin olive oil fit for consumption as it is * includes:

i) Extra virgin olive oil: virgin olive oil which has a free acidity, expressed as oleic acid, of not more than 1 gram per 100 grams, and the organoleptic characteristics of which correspond to those fixed for this category in this standard.

^{*} Oil which may be referred to as "natural".

- ii) <u>Virgin olive oil</u> (the qualifier "fine" may be used at the production and wholesale stage): virgin olive oil which has a free acidity, expressed as oleic acid, of not more than 2 grams per 100 grams and the organoleptic characteristics of which correspond to those fixed for this category in this standard.
- iii) Ordinary virgin olive oil: virgin olive oil which has a free acidity, expressed as oleic acid, of not more than 3.3 grams per 100 grams and the organoleptic characteristics of which correspond to those fixed for this category in this standard.
- **2.1.1.2.** <u>Virgin olive oil not fit for consumption as it is</u>, designated <u>lampante virgin olive oil</u>, is virgin olive oil which has a free acidity, expressed as oleic acid, of more than 3.3 grams per 100 grams and/or the organoleptic characteristics of which correspond to those fixed for this category in this standard. It is intended for refining or for technical purposes.
- **2.1.2.** Refined olive oil is the olive oil obtained from virgin olive oils by refining methods which do not lead to alterations in the initial glyceridic structure.
- **2.1.3.** Olive oil is the oil consisting of a blend of refined olive oil and virgin olive oil fit for consumption as it is.
- **2.2.** Olive-pomace oil is the oil obtained by treating olive pomace with solvents, to the exclusion of oils obtained by re-esterification processes and of any mixture with oils of other kinds. It is marketed in accordance with the following designations and definitions:
- **2.2.1.** Crude olive-pomace oil is olive-pomace oil intended for refining with a view to its use in food for human consumption, or intended for technical purposes.
- **2.2.2.** <u>Refined olive-pomace oil</u> is the oil obtained from crude olive-pomace oil by refining methods which do not lead to alterations in the initial glyceridic structure.
- **2.2.3.** Olive-pomace oil is the oil comprising the blend of refined olive-pomace oil and virgin olive oil fit for consumption as it is. In no case shall this blend be called "olive oil".

Olive oils and

3. PURITY CRITERIA

The identity characteristics comprising the purity criteria shall be applicable to olive oil and olive-pomace oil.

The limits established for each criterion include the precision values of the attendant recommended method.

3.1. Sterol composition (% of total sterols)

	olive-pomace oils
 Cholesterol Brassicasterol Campesterol Stigmasterol Delta-7-stigmastenol Beta-sitosterol + 	
delta-5avenasterol + delta-5-23-stigmastadienol + clerosterol + sitostanol + delta 5-24-stigmastadienol 3.2. Total sterols content (mg/kg)	<u>></u> 93.0
 Virgin olive oils Refined olive oil Olive oil Crude olive-pomace oil Refined olive-pomace oil Olive-pomace oil) ≥ 1000) ≥ 2500 ≥ 1800 ≥ 1600

^{*} Limit raised to \leq 0.2 for olive-pomace oils.

3.3. <u>Fatty acid composition using gas-liquid chromatography</u> (% m/m of methyl esters):

-	Myristic acid	≤ 0.05
-	Palmitic acid	7.5 - 20.0
-	Palmitoleic acid	0.3 - 3.5
-	Heptadecanoic acid	<u><</u> 0.3
-	Heptadecenoic acid	≤ 0.3
-	Stearic acid	0.5 - 5.0
-	Oleic acid	55.0 - 83.0
-	Linoleic acid	3.5 - 21.0
-	Linolenic acid	<u><</u> 1.0
_	Arachidic acid	<u><</u> 0.6
-	Gadoleic acid (eicosenoic)	<u><</u> 0.4
-	Behenic acid	<u><</u> 0.2*
-	Lignoceric acid	<u><</u> 0.2

3.4. Saturated fatty acid content in the 2-position in the triglycerides: The maximum acceptable level is the sum of the palmitic and stearic acids:

-	Virgin olive oils	$\leq 1.5\%$
-	Refined olive oil	<u><</u> 1.8%
-	Olive oil	<u><</u> 1.8%
-	Crude olive-pomace oil	$\leq 2.2\%$
-	Refined olive-pomace oil	$\leq 2.2\%$
-	Olive-pomace oil	$\leq 2.2\%$

3.5. <u>Unsaponifiable matter</u>

-	Olive oils	\leq 15 g/kg
-	Olive-pomace oils	≤ 30 g/kg

^{*} Limit raised to ≤ 0.3 for olive-pomace oils.

3.6. Detection of olive-pomace oil

	Lampante virgin olive oil	Edible virgin oils	Refined olive oil	Olive oil	
Waxes mg/kg C40+C42+C44+C46	<u><</u> 350	<u><</u> 250	<u><</u> 350	<u>≤</u> 350	
Erythrodiol + uvaol/ total sterols %	<u>≤</u> 4.5	<u><</u> 4.5	<u><</u> 4.5	<u><</u> 4.5	

3.7. <u>Detection of seed oils</u>

Maximum difference between the real and theoretical ECN 42 triglyceride content:

_	Edible virgin olive oils	0.2
-	Refined olive oil	0.3
-	Olive oil	0.3
-	Lampante virgin olive oil	0.3
-	Refined olive-pomace oil	0.5
-	Olive-pomace oil	0.5
-	Crude olive-pomace oil	0.6

3.8. Detection of refined vegetable oils

Detection of Termed vegetation one	Stigmastadienes ppm	R1
 Edible virgin olive oils Lampante virgin olive oil Refined olive oil Olive oil Crude olive-pomace oil Refined olive pomace oil Olive-pomace oil 		$\begin{array}{c} \ge 12 \\ \ge 12 \\ \text{no limit} \\ \ge 10 \\ \ge 10 \end{array}$

The R1 ratio = stigmasta-3,5-diene is to be applied to oils whose campesta-3,5-diene

stigmastadiene content is greater than 4 ppm.

3.9. Trans fatty acid content

	C18:1 T	C18:2 T + C18:3 T %
 Edible virgin olive oils Lampante virgin olive oil Refined olive oil Olive oil Crude olive-pomace oil Refined olive-pomace oil Olive-pomace oil 	$ \leq 0.05 $ $ \leq 0.10 $ $ \leq 0.20 $ $ \leq 0.20 $ $ \leq 0.20 $ $ \leq 0.40 $ $ \leq 0.40 $	

^{*} Provisional limits

4. QUALITY CRITERIA

The limits established for each criterion and designation include the precision values of the attendant recommended method

Olive- pomace oil	poob		light,	yellow to green	limpid	^ 1 1.	> 1		< 1.70	< 0.18
Refined olive-pomace oil	acceptable		light,	yellow to brownish yellow	limpid	۸۱ 8.	۸۱ ص		≥ 2.00	≥ 0.20
Crude olive- pomace oil						no limit	no limit	-		
Olive	poob		light,	yellow to green	limpid	s:1.5	^1 5		< 0.90	≤ 0.15
Refined olive oil	acceptable			light yellow	limpid	۸۱ 6.	ιο V I		<u>< 1.10</u>	≥ 0.16
Lampante virgin olive oil *			Me > 6.0			۷ ښ	no limit		no limit**	
Virgin Ordinary Lampante olive oilve oil oil oil oil oil oil oil *			2.5 <me 6.0***<="" <="" td=""><td></td><td></td><td>۶. د. د.</td><td>. 20 . 20</td><td></td><td>< 0.30 **</td><td>≥ 0.01</td></me>			۶. د. د.	. 20 . 20		< 0.30 **	≥ 0.01
Virgin olive oil			0 < Me < 2.5 Me > 0			1 2.0	- 50 - 50		≥ 0.25	≥ 0.01
Extra virgin olive oil			Me = 0 Me > 0			1.0	<u> </u>		< 0.25	≥ 0.01
	4.1 <u>Organoleptic</u> <u>characteristics</u> - odour and taste	- odour and taste (on a continuous scale):	. median of defect . median of the fruity attribute	- colour	- aspect at 20°C for 24 hours	4.2. Free acidity % m/m expressed in oleic acid	4.3. <u>Peroxide value</u> in milleq. Peroxide oxygen per kg/oil	4.4. Absorbency in ultra-violet (K ^{1%})	- 270 nm	- ∆ K

^{*} It is not obligatory for the criteria in 4.1, 4,2 and 4.3 to be concurrent; one is sufficient.

** After passage of the sample through activated alumina, absorbency at 270 nm shall be equal to or less than 0.11.

*** Or when the median of the defect is less than or equal to 2.5 and the median of the fruity attribute is equal to 0.

5. FOOD ADDITIVES

5.1. Virgin olive oils and crude olive-pomace oil:

none permitted.

5.2. Refined olive oil, olive oil, refined olive-pomace oil and olive-pomace oil: alpha-tocopherol permitted to restore natural tocopherol lost in the refining process.

Maximum level: 200 mg/kg of total alpha-tocopherol in the final product.

COI/T.15/NC no. 2/Rev. 8 page 9

6. CONTAMINANTS

Olive- pomace oil	۸۱ ۲۰	0.05	ī	۸۱۸۱ 8.0 1.0		۸۱ ۲.0	< 0.2	
Refined olive-pomace oil	× 0.1	0.05	T	6 0.0 0.0 0.1		۸۱ ۲.0	< 0.2	
Crude olive- pomace oil	۸۱ د :		> 120°C					
Olive	1.0 ^1	< 0.05	1	A1 A1 0.1.0		۱۸ ۲.0	< 0.2	
Refined olive oil	۸۱ د.	< 0.05	,	A 3.0		≥0.1	< 0.2	
Lampante virgin olive	o o o o o o o o o o o o o o o o o o o	< 0.2	Ť	3.01.01.01.0				
Ordinary virgin olive	S. 0. 5. 1	۸۱ د.	,	AI AI 8.0 1.0		۱۵ ما ۲.0 ما		
Virgin olive oil	< 0.2	۸۱ ۲.0	ı	6 0.0 0.1.0	2.	< 0.1	≥ 0.2	
Extra virgin olive	> 0.2	× 0 × 1	ı	A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1 A 1		۸۱ ۲.	< 0.2	
	6.1 <u>Moisture and volatile matter</u> (% m/m)	6.2. <u>Insoluble</u> <u>impurities</u> (% m/m) in light petroleum	6.3. Flash point	6.4 <u>. Trace metals</u> mg/kg Iron Copper	6.5. <u>Halogenated</u> solvents	Each solvent detected Mg/kg	Sum of solvents detected, mg/kg	

7. HYGIENE

It is recommended that the products intended for human consumption covered by the provisions of this standard be prepared in accordance with the appropriate sections of the General Principles of Food Hygiene recommended by the Codex Alimentarius Commission (CAC/RCP 1 - 1969, Rev. 2 - 1985).

8. PACKING

Olive oils and olive-pomace oils intended for international trade shall be packed in containers complying with the General Principles of Food Hygiene recommended by the Codex Alimentarius Commission (CAC/RCP 1 - 1969, Rev. 2 - 1985).

The containers used may be:

- **8.1.** <u>tanks</u>, <u>containers</u>, <u>vats</u>, which permit the transportation in bulk of olive oils and olive-pomace oils;
- **8.2.** metal drums, in good condition, hermetically-sealed, which should be internally covered with a suitable varnish;
- **8.3.** metal tins and cans, lithographed, new, hermetically-sealed, which should be internally covered with a suitable varnish;
- **8.4.** <u>demi-johns, glass bottles</u> or bottles made of suitable macromolecular material.

9. <u>CONTAINER FILLING TOLERANCE</u>

The volume occupied by the contents shall under no circumstances be less than 90% of the capacity of the container, except in the case of tin containers with a capacity of, or less than, 1 litre in which the volume occupied shall under no circumstances be less than 80% of the capacity of the container; this capacity is equal to the volume of distilled water at 20°C which the container can hold when full.

10. LABELLING

In addition to sections 2, 3, 7 and 8 of the Codex General Standard for the Labelling of Pre-packaged Foods (CODEX STAN 1-1985, Rev.1 - 1991) and the guidelines applying to food not intended for direct sale to consumers, the specific provisions providing the following information shall be applied:

10.1. On containers intended for direct sale to consumers

10.1.1. Name of the product

The labelling on each container shall indicate the specific designation of the product contained, complying in every way with the relevant provisions of this standard.

10.1.1.1. Designations of olive oil:

- Extra virgin olive oil*
- Virgin olive oil*
- Ordinary virgin olive oil*
- Refined olive oil
- Olive oil**

10.1.1.2. Designations of olive-pomace oil:

- Refined olive-pomace oil
- Olive-pomace oil.

10.1.2. Free acidity of the oil

The free acidity of the oil shall be declared on the label and expressed in terms of oleic acid (percentage m/m or degrees).

10.1.3. Net contents

The net contents shall be declared by weight or volume in the metric system ("Système International" units).

10.1.4. Name and address

The name and address of the manufacturer, packer, distributor, importer, exporter or seller shall be declared.

^{*} Oil which may likewise be referred to as "natural".

^{**} The terms "pure" or "100% pure" may figure on the label as a specification of the product.

10.1.5. Country of origin

The name of the country of origin shall be declared. When the product undergoes processing or re-packing, including in small containers, in a second country, the country in which the processing was performed shall be considered the country of origin for the purposes of labelling.

10.1.6. <u>Indications of source and appellations of origin</u>

10.1.6.1. Indications of source

The labels of virgin olive oils may indicate their source (country, region or locality) when they have been empowered to do so by their country of origin and when such virgin olive oils have been produced, packed and originate exclusively in the country, region or locality mentioned.

The labels for blends of refined olive oil and virgin olive oil may only indicate the source of the exporting country.

10.1.6.2. Appellations of origin

The labels of extra virgin olive oils may indicate their appellation of origin (country, region or locality) when they have been awarded such an appellation, in accordance with the terms provided under the regulations of their country of origin and when such extra virgin olive oil has been produced, packed and originates exclusively in the country, region or locality mentioned.

10.1.7. Lot identification

Each container shall be embossed or otherwise permanently marked in code or in clear to identify the producing factory and the lot.

10.1.8. Date marking and storage conditions

10.1.8.1. Date of packing

The date of packing shall be declared by the month and year in uncoded numerical sequence.

The month may be indicated by letters in those countries where such use will not confuse the consumer; if the month is December, the expression "end (stated year)" may be used as an alternative.

10.1.8.2. Date of minimum durability

In the case of pre-packaged products intended for the end consumer, the date of minimum durability (preceded by the words "best before end") shall be declared by the month and year in uncoded numerical sequence. The month may be indicated by letters in those countries where such use will not confuse the consumer; if the shelf life of the product is valid to December, the expression "end (stated year)" may be used as an alternative.

The period of durability shall not exceed 12 months after the date of packing. It may, however, be extended to 18 months for oils packed in metal containers.

10.1.8.3. Storage instructions

Any special conditions for storage shall be declared on the label if the validity of the date of minimum durability depends thereon.

10.2. On forwarding packs of oils intended for human consumption

In addition to the details noted under section 10.1., the following inscription shall appear:

- number and type of containers held in pack.

10.3. On containers allowing the transportation in bulk of olive oils and olive-pomace oils

The labelling on each container shall include:

10.3.1. Name of the product

The name shall indicate the specific designation of the product contained, complying in every way with the provisions of this standard.

10.3.2. Net contents

The net contents shall be declared by weight or volume in the metric system ("Système International" units).

10.3.3. Name and address

The name and address of the manufacturer, distributor or exporter shall be declared.

10.3.4. Country of origin

The name of the exporting country shall be declared.

11. METHODS OF ANALYSIS AND SAMPLING

The methods of analysis and sampling given below are international referee methods. The latest version of these methods should be used.

11.1. Sampling

According to ISO method 5555, "Animal and vegetable fats and oils - Sampling".

11.2. Preparation of the test sample

According to ISO method 661, "Animal and vegetable fats and oils - Preparation of the test sample".

11.3. Determination of the fatty acid composition

According to the capillary column ISO method 5508, "Analysis by Gas Chromatography of Methyl Esters of Fatty Acids" and ISO 5509, "Preparation of Fatty Acid Methyl Esters".

11.4. Determination of the unsaponifiable matter

According to IUPAC method no. 2.401, "Determination of the unsaponifiable matter", using diethyl ether, or ISO 3596 - 1.

The results are expressed in g/unsaponifiable matter per kg/oil.

11.5. Detection of olive-pomace oil

According to the following methods:

- COI/T.20/Doc. no. 18, "Determination of Wax Content by Capillary-Column Gas Liquid Chromatography".
- IUPAC no. 2.431, "Determination of the Erythrodiol Content". Capillary columns are recommended.

11.6. Detection of seed oils

According to the method COI/T.20/Doc. no. 20, "Determination of the difference between real and theoretical ECN 42 triglyceride content".

The prior purification of the oils should be performed according to IUPAC method no. 2.507, "Determination of the polar compounds in frying fats".

11.7. Detection of refined vegetable oils

According to the following methods:

- COI/T.20/Doc. no. 11, "Determination of Stigmastadienes in Vegetable Oils";
- COI/T.20/Doc. no. 16, "Determination of Sterenes in refined Vegetable Oils".

11.8. Determination of the trans fatty acid content

According to the method COI/T.20/Doc. no. 17, "Determination of Trans Unsaturated Fatty Acids by Capillary Column Gas Chromatography.

11.9. Determination of the sterol composition and total sterols content

According to the method COI/T.20/Doc. no. 10, "Determination of the Composition and Content of Sterols by Capillary-Column Gas Chromatography".

11.10. Determination of the fatty acids in the 2-position in the triglycerides

According to IUPAC method no. 2.210, "Determination of the Fatty Acids in the 2-Position in the Triglycerides of Oils and Fats", or ISO 6800.

11.11. Determination of the organoleptic characteristics

According to the method COI/T.20/Doc. no. 15, "Organoleptic Assessment of Virgin Olive Oil".

11.12. Determination of the free acidity

According to ISO method 660, "Determination of Acid Value and Acidity".

11.13. Determination of the peroxide value

According to IUPAC method no. 2.501, "Determination of the Peroxide Value (P.V.)", or to ISO 3960.

11.14. Determination of the absorbency in ultra-violet

According to the method COI/T.20/Doc. no. 19, "Spectrophotometric investigation in the ultraviolet".

11.15. Determination of the alpha-tocopherol

According to IUPAC method no. 2.432, "Identification and Determination of Tocopherols".

11.16. Determination of the moisture and volatile matter

According to IUPAC method no. 2.601, "Determination of the Moisture and Volatile Matter", or to ISO 662.

11.17. Determination of the insoluble impurities in light petroleum

According to IUPAC method no. 2.604, "Determination of the Insoluble Impurities", or to ISO 663.

11.18. Determination of the flash point

According to the FOSFA International method.

11.19. Detection of trace metals

According to IUPAC method no. 2.631, "Determination of Copper, Iron and Nickel by Direct Graphite Furnace Atomic Absorption Spectrometry", or ISO 8294.

11.20. Detection of traces of halogenated solvents

According to the method COI/T.20/Doc. no. 8/Corr. 1, "Determination of Tetrachloroethylene in Olive Oils by Gas-Liquid Chromatography".