

DETERMINATION OF THE DIFFERENCE BETWEEN ACTUAL AND THEORETICAL CONTENT OF TRIACYGLYCEROLS WITH ECN 42

1. Scope

Determination of the difference between the theoretical value of triacylglycerols (TAGs) with an equivalent carbon number of 42 (ECN 42_{theoretical}) calculated from the fatty acid composition, and the analytical results (ECN 42_{HPLC}) obtained by determination in the oil by high performance liquid chromatography.

2. Field of application

The standard is applicable to olive oils. The method is applicable to the detection of the presence of small amounts of seed oils (rich in linoleic acid) in every class of olive oils.

3. Principle

The content of triacylglycerols with ECN 42 determined by HPLC analysis and the theoretical content of triacylglycerols with ECN 42 (calculated on the basis of GLC determination of fatty acid composition) correspond within a certain limit for genuine olive oils. A difference larger than the values adopted for each type of oil points out that the oil contains seed oils.

4. Method

The method for the calculation of the theoretical content of triacylglycerols with ECN 42 and of the difference with respect to the HPLC data is essentially by the co-ordination of analytical data obtained by means of other methods. It is possible to distinguish three phases: determination of fatty acid composition by capillary gas chromatography, calculation of theoretical composition of triacylglycerols with ECN 42, HPLC determination of ECN 42 triacylglycerols.

4.1. Apparatus

- 4.1.1 Round-bottomed flasks, 250 and 500 mL.
- 4.1.2 Beakers 100 mL.
- 4.1.3 Glass chromatographic column, 21 mm internal diameter, 450 mm length, with cock and normalised cone (female) at the top.
- 4.1.4 Separating funnels, 250 mL, with normalised cone (male) at the bottom, suitable for connection to the top of the column.
- 4.1.5 Glass rod, 600 mm length.
- 4.1.6 Glass funnel, 80 mm diameter.
- 4.1.7 Volumetric flasks, 50 mL.
- 4.1.8 Volumetric flasks, 20 mL.
- 4.1.9 Rotary evaporator.
- 4.1.10 High performance liquid chromatograph, allowing thermostatic control of column temperature.
- 4.1.11 Injection units for 10 µl delivery.
- 4.1.12 Detector: differential refractometer. The full scale sensitivity should be at least 10^{-4} units of refractive index.
- 4.1.13 Column: stainless steel tube 250 mm length x 4.5 mm internal diameter packed with 5 µm diameter particles of silica with 22 to 23% carbon in the form of octadecylsilane*.
- 4.1.13 Recorder and/or integrator.

*Examples: Lichrosorb (Merck) RP 18 Art 50333

Lichrosphere (Merck) 100 CH18 Art 50377 or equivalent

4.2. Reagents

The reagents should be of analytical purity. Elution solvents should be de-gassed, and may be recycled several times without effect on the separations.

- 4.2.1 Petroleum ether 40-60°C chromatographic grade.
- 4.2.2 Ethyl ether, peroxide-free, freshly distilled.
- 4.2.3 Elution solvent for purifying the oil by column chromatography mixture petroleum ether/ethyl ether 87/13 (v/v).
- 4.2.4 Silicagel, 70-230 mesh, type Merck 7734, with water content standardised at 5% (w/w/).

- 4.2.5 Glass wool.
- 4.2.6 Acetone for HPLC.
- 4.2.7 Acetonitrile for HPCL.
- 4.2.8 HPLC elution solvent: acetonitrile + acetone (proportions to be adjusted to obtain the desired separation; begin with 50:50 mixture).
- 4.2.9 Solubilisation solvent: acetone.
- 4.2.10 Reference triglycerides: commercial triglycerides (tripalmitin, triolein, etc.) may be used and the retention times then plotted in accordance with the equivalent carbon number, or alternatively reference chromatograms obtained from soya oil, mixture 30:70 soya oil - olive oil and pure olive oil (see notes 1 and 2 and figures 1, 2, 3, 4).

4.3. Sample preparation

As a number of interfering substances can give rise to false positive results, the sample must always be purified according to IUPAC method 2.507, used for the determination of polar compounds in frying fats.

4.3.1 Chromatographic column preparation

Fill the column (4.1.3.) with about 30 mL of elution solvent (4.2.3.), then introduce inside the column some glass wool (4.2.5.) pushing it to the bottom of the column by means of the glass rod (4.1.5.)

In a 100 mL beaker, suspend 25 g of silicagel (4.2.4.) in 80 mL of elution mixture (4.2.3.), then transfer it to the column by means of a glass funnel (4.1.6.).

To ensure the complete transfer of the silicagel to the column, wash the beaker with the elution mixture and transfer the washing portions to the column too.

Open the cock and let the solvent elute from the column until its level is about 1 cm over the silicagel.

4.3.2 Column chromatography

Weigh with the accuracy of 0.001 g, 2.5 ± 0.1 g of oil, previously filtered, homogenised and anhydriified, if necessary, in a 50 mL volumetric flask (4.1.7.). Dissolve it in about 20 mL of elution solvent (4.2.3.). If necessary, slightly heat it to make the dissolution easily. Cool at room temperature and adjust the volume with elution solvent.

By means of a volumetric pipette, introduce 20 mL of solution inside the column prepared according to 4.3.1., open the cock and let the solvent elute to the silicagel layer level.

Then elute with 150 mL of elution solvent (4.2.3.), adjusting the solvent rate at about 2 mL/min (150 mL will take about 60-70 minutes to pass through the column).

The eluate is recovered in a 250 mL round-bottomed flask (4.1.1.) previously tared in an oven and exactly weighed. Eliminate the solvent at reduced pressure (Rotavapor) and weigh the residue that will be used to prepare the solution for HPLC analysis and for methyl ester preparation.

The sample recovery from the column must be 90% at least for the extra virgin, virgin, ordinary, refined and olive oil categories, and a minimum of 80% for lampante and olive-pomace oils.

4.4. HPLC analysis

4.4.1 Preparation of the samples for chromatographic analysis

A 5% solution of the sample to be analysed is prepared by weighing 0.5 ± 0.001 g of the sample into a 10 ml graduated flask and making up to 10 ml with the solubilisation solvent (4.2.9.).

4.4.2 Procedure

Set up the chromatographic system. Pump elution solvent (4.2.8) at a rate of 1.5 ml/min to purge the entire system. Wait until a stable base line is obtained. Inject 10 μ l of the sample prepared as in 4.3.

4.4.3 Calculation and expression of results

Use the area normalisation method, i.e. assume that the sum of the areas of the peaks corresponding to TAGs from ECN 42 up to ECN 52 is equal to 100%.

Calculate the relative percentage of each triglyceride using the formula:

$$\% \text{ triglyceride} = \text{area of peak} \times 100 / \text{sum of peak areas.}$$

The results to be given with at least two decimal places.

See notes 1, 2, 3 and 4.

4.5. Calculation of triacylglycerols composition (moles %) from fatty acid composition data (area %)

4.5.1 Determination of fatty acid composition

Fatty acid composition is determined by ISO 5508 by means of a capillary column. The methyl esters are prepared according to ISO 5509 (sodium methylate alcohol solution).

4.5.2 Fatty acids for calculation

Glycerides are grouped by their Equivalent Carbon Number (ECN), taking into account the following equivalencies between ECN and fatty acids. Only fatty acids with 16 and 18 carbon atoms were taken into consideration, because only these are important for olive oil.

Fatty acid (FA)	Abbreviation	Molecular weight (MW)	ECN
Palmitic acid	P	256.4	16
Palmitoleic acid	Po	254.4	14
Stearic acid	S	284.5	18
Oleic acid	O	282.5	16
Linoleic acid	L	280.4	14
Linolenic acid	Ln	278.4	12

4.5.3 Conversion of area % into moles for all fatty acids

$$\begin{array}{l}
 \text{moles P} = \frac{\text{area \% P}}{\text{MW P}} \quad \text{moles S} = \frac{\text{area \% S}}{\text{MW S}} \quad \text{moles Po} = \frac{\text{area \% Po}}{\text{MW Po}} \\
 \text{moles O} = \frac{\text{area \% O}}{\text{MW O}} \quad \text{moles L} = \frac{\text{area \% L}}{\text{MW L}} \quad \text{moles Ln} = \frac{\text{area \% Ln}}{\text{MW Ln}}
 \end{array} \quad \left. \vphantom{\begin{array}{l} \text{moles P} \\ \text{moles O} \end{array}} \right\} (1)$$

4.5.4 Normalisation of fatty acids to 100%

$$\begin{array}{l}
 \text{moles \% P (1,2,3)} = \frac{\text{moles P} * 100}{\text{moles (P + S + Po + O + L + Ln)}} \\
 \text{moles \% S (1,2,3)} = \frac{\text{moles S} * 100}{\text{moles (P + S + Po + O + L + Ln)}} \\
 \text{moles \% Po (1,2,3)} = \frac{\text{moles Po} * 100}{\text{moles (P + S + Po + O + L + Ln)}} \\
 \text{moles \% O (1,2,3)} = \frac{\text{moles O} * 100}{\text{moles (P + S + Po + O + L + Ln)}} \\
 \text{moles \% L (1,2,3)} = \frac{\text{moles L} * 100}{\text{moles (P + S + Po + O + L + Ln)}} \\
 \text{moles \% Ln (1,2,3)} = \frac{\text{moles Ln} * 100}{\text{moles (P + S + Po + O + L + Ln)}}
 \end{array} \quad \left. \vphantom{\begin{array}{l} \text{moles \% P} \\ \text{moles \% S} \\ \text{moles \% Po} \\ \text{moles \% O} \\ \text{moles \% L} \\ \text{moles \% Ln} \end{array}} \right\} (2)$$

The result gives the percentage of each fatty acid in moles % in the overall (1, 2, 3-) position of the TAGs.

Then the sum of the saturated fatty acids P and S (SFA) and the unsaturated fatty acids Po, O, L and Ln (UFA) are calculated:

$$\begin{array}{l}
 \text{moles \% SFA} = \text{moles \% P} + \text{moles \% S} \\
 \text{moles \% UFA} = 100 - \text{moles \% SFA}
 \end{array} \quad \left. \vphantom{\begin{array}{l} \text{moles \% SFA} \\ \text{moles \% UFA} \end{array}} \right\} (3)$$

4.5.5 Calculation of the fatty acid composition in 2- and 1,3- positions of TAGs

The fatty acids are distributed to three pools as follows: two identical for 1- and 3-positions and one for 2-position, with different coefficients for the saturated (P and S) and unsaturated acids (Po, O, L and Ln).

4.5.5.1 Saturated fatty acids in 2-position [P(2) and S(2)]

$$\left. \begin{aligned} \text{moles \% P(2)} &= \text{moles \% P(1,2,3)} * 0.06 \\ \text{moles \% S(2)} &= \text{moles \% S(1,2,3)} * 0.06 \end{aligned} \right\} \quad (4)$$

4.5.5.2 Unsaturated fatty acids in 2-position [Po(2), O(2), L(2) and Ln(2)]:

$$\left. \begin{aligned} \text{moles \% Po(2)} &= \frac{\text{moles \% Po(1,2,3)}}{\text{moles \% UFA}} * (100 - \text{moles \% P(2)} - \text{moles \% S(2)}) \\ \text{moles \% O(2)} &= \frac{\text{moles \% O(1,2,3)}}{\text{moles \% UFA}} * (100 - \text{moles \% P(2)} - \text{moles \% S(2)}) \\ \text{moles \% L(2)} &= \frac{\text{moles \% L(1,2,3)}}{\text{moles \% UFA}} * (100 - \text{moles \% P(2)} - \text{moles \% S(2)}) \\ \text{moles \% Ln(2)} &= \frac{\text{moles \% Ln(1,2,3)}}{\text{moles \% UFA}} * (100 - \text{moles \% P(2)} - \text{moles \% S(2)}) \end{aligned} \right\} \quad (5)$$

4.5.5.3 Fatty acids in 1,3-positions [P(1,3), S(1,3), Po(1,3), O(1,3), L(1,3) and Ln(1,3)]:

$$\left. \begin{aligned} \text{moles \% P(1,3)} &= \frac{\text{moles \% P(1,2,3)} - \text{moles \% P(2)}}{2} + \text{moles \% P(1,2,3)} \\ \text{moles \% S(1,3)} &= \frac{\text{moles \% S(1,2,3)} - \text{moles \% S(2)}}{2} + \text{moles \% S(1,2,3)} \\ \text{moles \% Po(1,3)} &= \frac{\text{moles \% Po(1,2,3)} - \text{moles \% Po(2)}}{2} + \text{moles \% Po(1,2,3)} \end{aligned} \right\} \quad (6)$$

$$\begin{aligned}
 \text{moles \% O(1,3)} &= \frac{\text{moles \% O(1,2,3)} - \text{moles \% O(2)}}{2} + \text{moles \% O(1,2,3)} \\
 \text{moles \% L(1,3)} &= \frac{\text{moles \% L(1,2,3)} - \text{moles \% L(2)}}{2} + \text{moles \% L(1,2,3)} \\
 \text{moles \% Ln(1,3)} &= \frac{\text{moles \% Ln(1,2,3)} - \text{moles \% Ln(2)}}{2} + \text{moles \% Ln(1,2,3)}
 \end{aligned}
 \tag{6}$$

4.5.6 Calculation of triacylglycerols

4.5.6.1 TAGs with one fatty acid (AAA, here LLL, PoPoPo)

$$\text{moles \% AAA} = \frac{\text{moles \% A(1,3)} * \text{moles \% A(2)} * \text{moles \% A(1,3)}}{10,000}
 \tag{7}$$

4.5.6.2 TAGs with two fatty acids (AAB, here PoPoL, PoLL)

$$\begin{aligned}
 \text{moles \% AAB} &= \frac{\text{moles \% A(1,3)} * \text{moles \% A(2)} * \text{moles \% B(1,3)} * 2}{10,000} \\
 \text{moles \% ABA} &= \frac{\text{moles \% A(1,3)} * \text{moles \% B(2)} * \text{moles \% A(1,3)}}{10,000}
 \end{aligned}
 \tag{8}$$

4.5.6.3 TAGs with three different fatty acids (ABC, here OLLn, PLLn, PoOLn, PPOLn)

$$\begin{aligned}
 \text{moles \% ABC} &= \frac{\text{moles \% A(1,3)} * \text{moles \% B(2)} * \text{moles \% C(1,3)} * 2}{10,000} \\
 \text{moles \% BCA} &= \frac{\text{moles \% B(1,3)} * \text{moles \% C(2)} * \text{moles \% A(1,3)} * 2}{10,000} \\
 \text{moles \% CAB} &= \frac{\text{moles \% C(1,3)} * \text{moles \% A(2)} * \text{moles \% B(1,3)} * 2}{10,000}
 \end{aligned}
 \tag{9}$$

4.5.6.4 Triacylglycerols with ECN42

The triacylglycerols with ECN42 are calculated according to equations 7, 8 and 9 and are then given in order of expected elution in HPLC (normally only three peaks).

LLL

PoLL and the positional isomer LPoL

OLLn and the positional isomers OLnL and LnOL

PoPoL and the positional isomer PoLPo

PoOLn and the positional isomers OPoLn and OLnPo

PLLn and the positional isomers LLnP and LnPL

PoPoPo

SLnLn and the positional isomer LnSLn

PPoLn and the positional isomers PLnPo and PoPLn

The triacylglycerols with ECN42 are given by the sum of the nine triacylglycerols including their positional isomers. The results to be given to at least two decimal places.

5. Evaluation of the results

The calculated theoretical content and the content determined by the HPLC analysis are compared. If the difference HPLC data minus theoretical data is greater than the values stated for the appropriate oil category in the standard, the sample contains seed oil.

Results are given to one decimal figure.

6. EXAMPLE (The numbers refer to the sections in the text of the method)

- 4.5.1. Calculation of moles % fatty acids from GLC data (area %)

The following data are obtained for the fatty acid composition by GLC:

FA	P	S	Po	O	L	Ln
MW	256.4	284.5	254.4	282.5	280.4	278.4
Area %	10.0	3.0	1.0	75.0	10.0	1.0

- 4.5.3 Conversion of area % into moles for all fatty acids

$$\text{moles P} = \frac{10}{256.4} = 0.03900 \text{ moles P}$$

$$\text{moles S} = \frac{3}{284.5} = 0.01054 \text{ moles S}$$

$$\text{moles Po} = \frac{1}{254.4} = 0.00393 \text{ moles Po}$$

$$\text{moles O} = \frac{75}{282.5} = 0.26549 \text{ moles O}$$

$$\text{moles L} = \frac{10}{280.4} = 0.03566 \text{ moles L}$$

$$\text{moles Ln} = \frac{1}{278.4} = 0.003594 \text{ moles Ln}$$

See formula (1)

$$\text{Total} = 0.35822 \text{ moles TAGs}$$

- 4.5.4 Normalisation of fatty acids to 100%

$$\begin{aligned}
 \text{moles \% P(1,2,3)} &= \frac{0.03900 \text{ moles P} * 100}{0.35822 \text{ moles}} = 10.888\% \\
 \text{moles \% S(1,2,3)} &= \frac{0.01054 \text{ moles S} * 100}{0.35822 \text{ moles}} = 2.944\% \\
 \text{moles \% Po(1,2,3)} &= \frac{0.00393 \text{ moles Po} * 100}{0.35822 \text{ moles}} = 1.097\% \\
 \text{moles \% O(1,2,3)} &= \frac{0.26549 \text{ moles O} * 100}{0.35822 \text{ moles}} = 74.113\% \\
 \text{moles \% L(1,2,3)} &= \frac{0.03566 \text{ moles L} * 100}{0.35822 \text{ moles}} = 9.956\% \\
 \text{moles \% Ln(1,2,3)} &= \frac{0.00359 \text{ moles Ln} * 100}{0.35822 \text{ moles}} = 1.003\%
 \end{aligned}$$

} See formula (2)

$$\text{Total moles \%} = 100.0\%$$

Sum of the saturated and unsaturated fatty acids in the 1,2,3-position of TAGs:

$$\begin{aligned}
 \text{moles \% SFA} &= 10.888\% + 2.944\% = \mathbf{13,831\%} \\
 \text{moles \% UFA} &= 100.000\% - 13.831\% = \mathbf{86,169\%}
 \end{aligned}$$

} See formula (3)

- 4.5.5 Calculation of the fatty acid composition in 2- and 1,3-positions of the TAGs

- 4.5.5.1 Saturated fatty acids in 2-position [P(2) and S(2)]

$$\begin{aligned}
 \text{moles \% P(2)} &= 10.888\% * 0.06 = 0.653 \text{ moles \%} \\
 \text{moles \% S(2)} &= 2.944\% * 0.06 = 0.177 \text{ moles \%}
 \end{aligned}$$

} See formula (4)

- 4.5.5.2 Unsaturated fatty acids in 2-position [Po(1,3), O(1,3), L(1,3) and Ln(1,3)]

$$\text{moles \% Po(2)} = \frac{1.097\%}{86.169\%} * (100 - 0.659 - 0.177) = 1.263 \text{ moles \%}$$

$$\text{moles \% O(2)} = \frac{74.113\%}{86.169\%} * (100 - 0.659 - 0.177) = 85.295 \text{ moles \%}$$

$$\text{moles \% L(2)} = \frac{9.956\%}{86.169\%} * (100 - 0.659 - 0.177) = 11.458 \text{ moles \%}$$

$$\text{moles \% Ln(2)} = \frac{1.003\%}{86.169\%} * (100 - 0.659 - 0.177) = 1.154 \text{ moles \%}$$

See formula (5)

- 4.5.5.3 Fatty acids in 1,3-positions [P(1,3), S(1,3), Po(1,3), O(1,3), L(1,3) and Ln(1,3)]

$$\text{moles \% P(1,3)} = \frac{10.888 - 0.659}{2} + 10.888 = 16.005 \text{ moles \%}$$

$$\text{moles \% S(1,3)} = \frac{2.944 - 0.177}{2} + 2.944 = 4.327 \text{ moles \%}$$

$$\text{moles \% Po(1,3)} = \frac{1.097 - 1.263}{2} + 1.097 = 1.015 \text{ moles \%}$$

$$\text{moles \% O(1,3)} = \frac{74.113 - 85.295}{2} + 74.113 = 68.522 \text{ moles \%}$$

$$\text{moles \% L(1,3)} = \frac{9.956 - 11.458}{2} + 9.956 = 9.205 \text{ moles \%}$$

$$\text{moles \% Ln(1,3)} = \frac{1.003 - 1.154}{2} + 1.003 = 0.927 \text{ moles \%}$$

See formula (6)

- 4.5.6. Calculation of triacylglycerols

From the calculated fatty acid composition in sn-2- and sn-1,3-positions:

FA in	1,3-pos	2-pos
P	16.005%	0.653%
S	4.327%	0.177%
Po	1.015%	1.263%
O	68.522%	85.295%
L	9.205%	11.458%
Ln	0.927%	1.154%
Sum	100.0%	100.0%

the following triacylglycerols are calculated:

LLL

PoPoPo

PoLL with 1 positional isomer

SLnLn with 1 positional isomer

PoPoL with 1 positional isomer

PPoLn with 2 positional isomers

OLLn with 2 positional isomers

PLLn with 2 positional isomers

PoOLn with 2 positional isomers

- 4.5.6.1 TAGs with one fatty acid (LLL, PoPoPo)

See formula (7)

$$\text{mol\%LLL} = \frac{9.205\% * 11.458\% * 9.205\%}{10,000} = \underline{\underline{0.09708 \text{ mol LLL}}}$$

$$\text{mol\%PoPoPo} = \frac{1.15\% * 1.263\% * 1.015\%}{10,000} = \underline{\underline{0.00013 \text{ mol PoPoPo}}}$$

- 4.5.6.2 TAGs with two fatty acids (PoLL, SLnLn, PoPoL)

See formula (8)

$$\text{mol\%PoLL+LLPo} = \frac{1.015\% * 11.458\% * 9.205\% * 2}{10,000} = 0.02141$$

$$\text{mol\%LPoL} = \frac{9.205\% * 1.263\% * 9.205\%}{10,000} = 0.01070$$

0.03211 mol PoLL

$$\text{mol\%SLnLn+LnLnS} = \frac{4.327\% * 1.154\% * 0.927\% * 2}{10,000} = 0.00093$$

$$\text{mol\%LnSLn} = \frac{0.927\% * 0.177\% * 0.927\%}{10,000} = 0.00002$$

0.00095 mol SLnLn

$$\text{mol\%PoPoL+LPoPo} = \frac{1.015\% * 1.263\% * 9.205\% * 2}{10,000} = 0.00236$$

$$\text{mol\%PoLPo} = \frac{1.015\% * 11.458\% * 1.015\%}{10,000} = 0.00118$$

0.00354 mol PoPoL

- 4.5.6.3 TAGs with three different fatty acids (PoPLn, OLLn, PLLn, PoOLn)

See formula (9)

$$\text{mol\%PPoLn} = \frac{16.005\% * 1.263\% * 0.927\% * 2}{10,000} = 0.00375$$

$$\text{mol\%LnPPo} = \frac{0.927\% * 0.653\% * 1.015\% * 2}{10,000} = 0.00012$$

$$\text{mol\%PoLnP} = \frac{1.015\% * 1.154\% * 16.005\% * 2}{10,000} = 0.00375$$

0.00762 mol PPoLn

$$\text{mol\%OLLn} = \frac{68.522\% * 11.458\% * 0.927\% * 2}{10,000} = 0.14577$$

$$\text{mol\%LnOL} = \frac{0.927\% * 85.295\% * 9.205\% * 2}{10,000} = 0.14577$$

$$\text{mol\%LLnO} = \frac{9.205\% * 1.154\% * 68.522\% * 2}{10,000} = 0.14577$$

0.43671 mol OLLn

$$\text{mol\%PLLn} = \frac{16.005\% * 11.458\% * 0.927\% * 2}{10,000} = 0.03400$$

$$\text{mol\%LnPL} = \frac{0.927\% * 0.653\% * 9.205\% * 2}{10,000} = 0.00111$$

$$\text{mol\%LLnP} = \frac{9.205\% * 1.154\% * 16.005\% * 2}{10,000} = 0.03400$$

0.06911 mol PLLn

$$\text{mol\%PoOLn} = \frac{1.015\% * 85.295\% * 0.927\% * 2}{10,000} = 0.01605$$

$$\text{mol\%LnPoO} = \frac{0.927\% * 1.263\% * 68.522\% * 2}{10,000} = 0.01605$$

$$\text{mol\%OLnPo} = \frac{68.522\% * 1.154\% * 1.015\% * 2}{10,000} = 0.01605$$

0.04815 mol PoOLn

ECN42 = 0.69540 mol TAGs

Note 1: The elution order can be determined by calculating the equivalent carbon numbers, often defined by the relation $ECN = CN - 2n$, where CN is the carbon number and n is the number of double bonds; it can be calculated more precisely by taking into account the origin of the double bond. If n_o , n_l and n_{ln} are the numbers of double bonds attributed to oleic, linoleic and linolenic acids respectively, the equivalent carbon number can be calculated by means of the relation of the formula:

$$EN = CN - d_o n_o - d_l n_l - d_{ln} n_{ln}$$

where the coefficient d_o , d_l and d_{ln} can be calculated by means of the reference triglycerides. Under the conditions specified in this method, the relation obtained will be close to:

$$ECN = CN - (2.60 n_o) - (2.35 n_l) - (2.17 n_{ln})$$

Note 2: With several reference triglycerides, it is also possible to calculate the resolution with respect to triolein:

$$\alpha = RT^1 / RT \text{ triolein}$$

by use of the reduced retention time $RT^1 = RT - RT \text{ solvent}$

The graph of $\log \alpha$ against f (number of double bonds) enables the retention values to be determined for all the triglycerides of fatty acids contained in the reference triglycerides - see Figure 1.

Note 3: The efficiency of the column should permit clear separation of the peak of trilinoein from the peaks of the triglycerides with an adjacent RT. The elution is carried out up to ECN 52 peak.

Note 4: A correct measure of the areas of all peaks of interest for the present determination is ensured if the second peak corresponding to ECN 50 is 50% of full scale of the recorder.

Figure 1: Graph of $\log \alpha$ against f (number of double bonds)

Number of double bonds

La: lauric acid; My: myric acid; P: palmitic acid; St: stearic acid; O: oleic acid; L: linoleic acid; Ln: linolenic acid

Figure 2: Soya oil

Figure 3: Soya oil/Olive oil 30/70

Figure 4: Olive oil

—

COMPUTER PROGRAM

PRECISION VALUES OF THE METHOD

1. Analysis of the collaborative test results

The precision values of the method are given in the table overleaf.

Nineteen laboratories holding IOOC recognition at the time took part in the collaborative test arranged by the Executive Secretariat in 1999. The laboratories were from eight countries.

The test was performed on five samples:

- A: extra virgin olive oil
- B: virgin olive oil + refined sunflower oil
- C: virgin olive oil + refined olive-pomace oil
- D: virgin olive oil + refined soybean oil + refined sunflower oil
- E: refined olive oil + refined olive-pomace oil + refined soybean oil + lampante virgin olive oil

The results of the collaborative test organised by the IOOC Executive Secretariat have been statistically processed according to the rules laid down in the international standards ISO 5725 **Accuracy (trueness and precision) of measurement methods and results**. Outliers were examined by applying Cochran's and Grubbs' test to the laboratory results for each determination (replicates a and b) and each sample.

The table lists :

- n** number of participating laboratories
- outliers** number of laboratories with outlying values
- mean** mean of the accepted results
- r** value below which the absolute difference between two single independent test results obtained with the same method on identical test material in the same laboratory by the same operator using the same equipment within short intervals of time may be expected to lie with a probability of 95%
- S_r** Repeatability standard deviation

RDS_r (%) Repeatability coefficient of variation ($S_r \times 100/\text{mean}$)

R value below which the absolute difference between two single test results obtained with the same method on identical test material in different laboratories with different operators using different equipment may be expected to lie with a probability of 95%

S_R Reproducibility standard deviation

RDS_R (%) Reproducibility coefficient of variation ($S_R \times 100/\text{mean}$)

Difference between actual and theoretical content of triglyceride with ECN 42

	A	B	C	D	E
n	19	19	19	19	19
outliers	1	0	0	0	3
mean	0.04	1.66	0.04	0.18	0.82
r	0.08	0.12	0.09	0.11	0.11
S_r	0.02	0.04	0.03	0.04	0.041
RSD_r (%)	82.24 _(not sig.)	2.77	76.11 _(not sig.)	22.51	5.07
R	0.12	0.25	0.16	0.22	0.24
S_R	0.05	0.09	0.05	0.08	0.08
RSD_R(%)	127.56 _(not sig.)	5.42	132.17 _(not sig.)	46.19	10.85

2. Normative references

ISO 5725-1: 1994 Accuracy (trueness and precision) of measurement methods and results – Part 1: General principles and definitions

ISO 5725-2: 1994 Accuracy (trueness and precision) of measurement methods and results – Part 2: Basic method for the determination of the repeatability and reproducibility of a standard measurement method

ISO 5725-5: 1994 Accuracy (trueness and precision) of measurement methods and results – Part 5: Alternative methods for the determination of the precision of a standard measurement method

ISO 5725-6: 1994 Accuracy (trueness and precision) of measurement methods and results – Part 6: Use in practice of accuracy values